

Request to remove head lice

There has been an outbreak of head lice. Head lice may appear as 'white specks that resembles dandruff,' 'itchiness on the child's head,' or 'frequent head scratching.' If your child is suspected of being infected, there is no need to cut their hair. Use lice removal medication and a fine-toothed comb, and follow these steps for removal:

Removing head lice using Sumithrin Shampoo

1 Wet the hair first. Apply the medication throughout the hair, and lather it as you would with regular shampoo.



2 Wrap a towel around the head and wait for 5 minutes.



3 Rinse out the medication and comb through the hair using a fine-toothed comb. Repeat this every three days for a total of 3-4 treatments.



Take note of these points!

- After rinsing out the medication, you may use regular shampoo and conditioner.
- Do not use the medication daily, either for prevention or to speed up their removal.
- Do not sprinkle Sumithrin Powder directly on pillowcases or sheets.

Remove head lice and eggs using a fine-toothed lice comb


Please use a special lice comb with a fine structure. Some combs have fine, needle-like teeth and are even designed in a screw-like pattern.

1 First, comb with your usual comb.

2 Next, divide the hair into small sections, hold the hair at the roots, and comb down to the ends using the lice removal comb. To remove the eggs physically, work slowly and gently to pull the eggs off the hair strands.



3 Soak the comb in a basin of hot water, and use a toothpick or similar tool to remove any eggs and lice caught in the comb's teeth.



4 After use, wash the comb thoroughly, dry it, and store it properly.

It is not necessary to suspend students infected with head lice from school; however, prompt and appropriate treatment is required.